

Drip Information

Effective: 03/01/2021

Supersedes:
N/A

DOSING AND ADMINISTRATION

MEDICATION	ADULT	PEDIATRIC (Refer to Handtevy)
<u>Norepinephrine</u>	5-15 mcg/min, titrated to SBP >90 or MAP >65.	0.05-0.1 mcg/kg/min titrated Q5 to age appropriate SBP or MAP Max dose 2 mcg/kg/min OR 30 mcg/min
<u>Amiodarone</u>	<u>Bolus:</u> 150mg over 10min IV/IO (100gtts/min. 150mg in 100cc of NS with 10gtts set)	<u>Bolus:</u> 150mg over 10min IV/IO (100gtts/min. 150mg in 100cc of NS with 10gtts set)
	<u>Maintenance Infusion:</u> 1mg/min IV/IO (40gtts/min. 150mg in 100cc NS with 60gtts set)	---
<u>Epinephrine</u>	2-10 mcg/min infusion (10-50 gtts/min, 1mg 1:1,000 in 500 ML NS with 10 drop set)	---
<u>Hydroxocobalamin</u>	5 g IV/IO over 15 minutes <u>In its own IV/IO line</u> (Run wide open with included drop set)	5 g IV/IO over 15 minutes <u>In its own IV/IO line</u> (Run wide open with included drop set)
<u>Magnesium Sulfate</u>	<u>Breathing Difficulty:</u> 1 g IV/IO over 20 minutes (60gtts/min in 100cc NS w/10gtt set)	25-50 mg/kg IV/IO over 20 minutes (60gtts/min in 100cc NS w/10gtt set)
	<u>Pre-Eclampsia / Eclampsia:</u> 4 g IV/IO over 20 minutes (60gtts/min in 100cc NS w/10gtt set)	